

## DETERMINING MODERN ASPECTS OF TREATMENT IN PATIENTS WITH HYPERACTIVE BLADDER SYNDROME

Shopulotov Shokhrukh Asliddinovich<sup>1</sup>, Shopulotova Zarina Abdumuminovna<sup>2</sup>, Tursunova Dilnura Akram kizi<sup>3</sup>

<sup>1</sup>Samarkand State Medical University, master degree department of Urology

<sup>2</sup>Samarkand State Medical University, assistant of department Obstetrics and gynecology №1

<sup>3</sup>Samarkand State Medical University, student of 430 group

<https://doi.org/10.5281/zenodo.10397474>

**Abstract.** *Overactive bladder (OBS) is a clinical syndrome involving urinary urgency with or without urge incontinence, usually in association with urinary frequency and nocturia (urination between falling asleep and waking up). To date, it has been established that the cause of frequent and urgent urination in most patients is detrusor hyperactivity. The problem of diagnosis and successful treatment of patients with OAB requires an integrated approach involving doctors of various specialties, primarily urologists, neurologists and neurosurgeons. Attention is drawn to the fact that in most cases it is not possible to determine the true cause of the development of OAB, which makes it necessary to continue research aimed at identifying the etiological factors of OAB.*

**Keywords:** *Overactive bladder (OAB) syndrome, urinary incontinence (UI), USE (ultrasound examination), stress urinary incontinence (SUI), urgent urinary incontinence (UUI), conservative treatment.*

**Relevance.** Overactive bladder syndrome (OBS) is a clinical syndrome with or without urgency incontinence, usually associated with frequent urination and nocturnal urination. To date, it has been established that detrusor hyperactivity is the cause of frequent and urgent urination in most patients. The problem of diagnosis and successful treatment of patients with OBS is a problem that requires the involvement of doctors of various specialties, primarily urologists, neurologists and neurosurgeons. In many cases, attention is paid to the fact that it is impossible to determine the real cause of the development of OBS, which requires the continuation of research aimed at determining the etiological factors of OBS [1-5, 7, 12].

Excessive activity of the bladder is not a life-threatening condition, but it has a sharp negative effect on its quality, leads to social adaptation and even disability [9, 15, 18].


Detrusor overactivity mimics or accompanies clinically stress urinary incontinence, making this form of incontinence difficult to detect and leading to inappropriate surgical correction of urinary retention, which in most cases can be avoided [4, 8]. The difficulty in diagnosing this condition is that detrusor hyperactivity is not felt by the patient, it is not accompanied by a desire to urinate, only symptoms of stress urinary incontinence are manifested [14, 17, 19]. Experience accumulated in recent years has shown that detrusor hyperactivity can be not only an independent cause of stress urinary incontinence, but can mimic the clinical presentation of this disease [11, 15, 16].

As can be seen from the above, bladder overactivity is an urgent problem in modern medicine. A detailed study of the etiological factors contributing to the development of OBS is required, as well as the development of measures for early diagnosis and prevention of OBS.

**The aim of the study.** Determine the methods of conservative treatment of patients with OBS.

**Materials and methods.** The study was based on the clinical and laboratory examination of 50 patients with hyperactive bladder syndrome who applied to the urology department of the Samarkand State Medical University in Samarkand and were hospitalized for treatment in 2021-2024.


The examined patients were divided into 2 groups depending on treatment methods (Fig. 1):



**Figure 1. Distribution of patients with urinary disorders by group**

The comparison group included 20 (40%) patients treated with conventional methods of diagnosis and treatment, and the main group included 30 (60%) patients treated with complex therapy.

The age of the examined patients was from 38 to 65 years. The mean age of patients in the comparison group was  $48.2 \pm 3.4$  years, and in the main group was  $49.03 \pm 4.12$  years, which is shown in Figure 2. by group, respectively.




**Figure 2. Distribution of the examined by age groups**

The distribution by gender was dominated by women (73.3% in the main group, 75% in the comparison group). It should be noted that the difference between women and men was significant, that is, men in all studies were aged 55 and over, while women had patients from 38 to 56 years and older.

All patients under observation underwent a thorough study of the somatic, urological, obstetrical and gynecological anamnesis, as well as a careful clinical examination, taking into account the age, the state of the reproductive system. General examination, external and internal urogynecological examination, general clinical analysis of blood and urine were performed. All laboratory analyzes were conducted in the laboratory department.

**Results and discussion.** In our study, according to the method of treatment, 50 patients were divided into 2 groups: main (n=30) and comparison (n=20) groups. Patients of the comparison group were treated conservatively with drugs. In addition to conservative drug therapy, physiotherapeutic method - BFB-training was applied to the main group of patients.

Drug therapy was prescribed according to the type of urinary incontinence. All patients were divided into Urgent Urinary Incontinence and Mixed Urinary Incontinence in Stress Overactive Bladder Syndrome (diagram 1).



**Diagram 1. Separation of patients of the main and comparison groups according to UI type**

For the treatment of urinary incontinence in hyperactive bladder syndrome, we used drugs of the M-cholinoblockers group. The mechanism of action of M-cholinoblockers is to block the muscarinic receptors of the urinary bladder, preventing the interaction of the mediator acetylcholine with them. Thus, the effect of acetylcholine on the detrusor is reduced or completely stopped, as a result of which its activity is reduced. In the treatment of detrusor hyperactivity, two types of M-cholinoblockers, differing in chemical structure, are used - tertiary and quaternary amines (oxybutynin, trospium chloride), which are the first-line drugs for hyperactive bladder

syndrome and ST. Oxybutynin was prescribed in a dose of 2.5 to 5 mg 3-4 times a day (maximum dose 20 mg/day).

The duration of the daily procedure varies between 15-20 minutes. The treatment course consists of 15 treatments. If necessary, repeated courses of BFB-therapy (2-3 times a year) can be conducted to enhance the clinical effect.

A total of 12 women with UI in OBS (5 comparison group and 7 main group) were prescribed beta-3-adrenomimetics with M-cholinoblockers during the first period of treatment. As a result, out of a total of 12 women, only 2 (3.33%; 5%) had a therapeutic effect from drug therapy and a positive result was obtained. When BFB was prescribed to 7 patients in the main group, positive dynamics were observed in all of them, and UI symptoms were completely eliminated in 3 patients.

Patients with the stress form of UI were prescribed antidepressants as a conservative treatment and the outcome was evaluated after 1 month. 7 people (35%) in the comparison group and 11 people (36.7%) in the main group had ST of this form.

At the same time as conservative treatment, BFB - training physiotherapeutic treatment method was applied to women of the main group. At this stage, the effectiveness of treatment was noted as positive in only 2 women (10%) in the comparison group, and in 8 women (26.7%) in the main group. At this stage, we can see that the effectiveness of complex treatment is several times higher than that of conventional treatment. Analysis of the obtained results showed that 17 (56.7%) patients in the main group did not have UI cases after complex treatment from OAB, this indicator was recorded in 4 (20%) patients in the comparison group, which is instead proves the effectiveness of complex therapy very reliably ( $P < 0.001$ ).

**Conclusion.** Analysis of the dynamics of the results of examinations of women with urinary incontinence by means of non-invasive special tests before and after treatment showed that after complex treatment 56.7% of patients did not have spontaneous urination during the tests, 43.3% of the main group patients did not fully recover, but reported positive results in terms of overall symptoms and a significant reduction in UI episodes. In 4 patients (20%) of the comparison group, spontaneous urinary excretion was not observed at all, and in the remaining 60%, the number of excretions decreased, but complete recovery was not observed, and in 20% of patients, there was no overall change before and after treatment.

As a result of the tests, all patients had the result of treatment. However, according to the results of the effectiveness, more patients of the main group had complete treatment compared to the comparison group. It is worth noting that during the course of treatment, they noted changes in positive dynamics from severe UI to mild UI, which led to a spontaneous reduction in the need for surgical treatment, as well as an improvement in the overall quality of life. will bring.

## REFERENCES

1. Amirzoda T. S., Asliddinovich S. S. НЕДЕРЖАНИЕ МОЧИ И НЕУДЕРЖАНИЕ МОЧИ: КАЧЕСТВО ЖИЗНИ ПАЦИЕНТОВ //JOURNAL OF BIOMEDICINE AND PRACTICE. – 2022. – Т. 7. – №. 5.
2. Мазо Е. Б., Кривобородов Г. Г. Гиперактивный мочевого пузыря //Consilium medicum. – 2003. – Т. 5. – №. 7. – С. 405-411

3. Неймарк А.И., Раздорская Мирослава Витальевна, Войтенко А.Н. Гиперактивный мочевого пузыря у женщин со стрессовым недержанием мочи // Андрология и генитальная хирургия. 2013. №1.
4. Сивков А. В., Ромих В. В. Фармакотерапия гиперактивного мочевого пузыря //Consilium medicum. – 2002. – Т. 4. – №. 7. – С. 348-355
5. Шопулотов Ш., Абсаматов Ш., Холдоров И. ГИПЕРАКТИВНОСТЬ ДЕТРУЗОРА: СОВРЕМЕННЫЕ ВОЗМОЖНОСТИ ТЕРАПИИ //Евразийский журнал медицинских и естественных наук. – 2022. – Т. 2. – №. 6. – С. 409-411.
6. Шопулотова З. А., Худоярова Д. Р. ХРОНИЧЕСКИЙ ПИЕЛОНЕФРИТ И БЕРЕМЕННОСТЬ: ВЛИЯНИЕ КОМОРБИДНОСТИ И ЕГО ПРОЯВЛЕНИЯ //ЖУРНАЛ РЕПРОДУКТИВНОГО ЗДОРОВЬЯ И УРО-НЕФРОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ. – 2023. – Т. 4. – №. 4.
7. Худоярова Д. Р., Уктамова Ю. У., Шопулотова З. А. ИННОВАЦИИ В КОМПЛЕКСНОМ ЛЕЧЕНИИ ХРОНИЧЕСКОГО ВОСПАЛЕНИЯ ПРИДАТКОВ МАТКИ //ЖУРНАЛ РЕПРОДУКТИВНОГО ЗДОРОВЬЯ И УРО-НЕФРОЛОГИЧЕСКИХ ИССЛЕДОВАНИЙ. – 2023. – Т. 4. – №. 4.
8. Shernazarov Farrukh ORGANIZATION OF DIGITALIZED MEDICINE AND HEALTH ACADEMY AND ITS SIGNIFICANCE IN MEDICINE // SAI. 2023. №Special Issue 8. URL: <https://cyberleninka.ru/article/n/organization-of-digitalized-medicine-and-health-academy-and-its-significance-in-medicine> (дата обращения: 20.11.2023).
9. Худоярова Д. Р. и др. Ведение беременных с преэклампсией тяжелой степени //Достижения науки и образования. – 2020. – №. 7 (61). – С. 29-30.
10. Худоёрова Д., Кобилова З., Шопулатов Ш. INFLUENCE OF CORONAVIRUS INFECTION ON CHILDREN AND THEIR MOMS //Журнал кардиореспираторных исследований. – 2020. – Т. 1. – С. 74-75.
11. Худоярова Д. Р. и др. Совершенствование методов естественного восстановления фертильности у женщин с бесплодием, обусловленным эндометриозом //Международный журнал" Journal of Biomedicine and Practice. – 2020. – №. 2. – С. 42.
12. Тохиров Х. и др. ХИРУРГИЧЕСКИЙ АБОРТ И ЕГО ПОСЛЕДСТВИЯ //Центральноазиатский журнал образования и инноваций. – 2023. – Т. 2. – №. 6 Part 4. – С. 21-23.
13. Hampel C, Weinhold D, Benken N et al. Prevalence and Natural History of Female Incontinence. Eur Urol 2007; 32 (Suppl. 2): 3–12
14. KHUDOYAROVA D. R. et al. RESTORING NATURAL FERTILITY IN WOMEN WITH ENDOMETRIOSIS //БИОМЕДИЦИНА ВА АМАЛИЁТ ЖУРНАЛИ. – С. 43.
15. Mukhsinov F. B. et al. Management of pregnant women with severe preeclampsia //Journal" Achievement of science and education. – 2020. – №. 7. – С. 61.
16. Shopulotov S., Isroilov O. GIPERAKTIV QOVUQ SINDROMINI DAVOLASH BO'YICHA YANGI IMKONIYATLAR //Бюллетень студентов нового Узбекистана. – 2023. – Т. 1. – №. 6 Part 2. – С. 106-109.
17. Shopulotov S. et al. SIYDIK TUTOLMASLIK MUAMMOSI //Центральноазиатский журнал образования и инноваций. – 2023. – Т. 1. – №. 6 Part 6. – С. 44-48.

18. Shopulotov S. et al. GIPERAKTIV QOVUQ SINDROMINI TASHXISLASHDA ZAMONAVIY YONDASHUVLAR //Молодые ученые. – 2023. – Т. 1. – №. 9. – С. 38-42.
19. Shopulotova Z. COMPARATIVE ANALYSIS OF CLINICAL CASES OF EXACERBATION OF CHRONIC PYELONEPHRITIS IN PREGNANT WOMEN //International Bulletin of Medical Sciences and Clinical Research. – 2023. – Т. 3. – №. 8. – С. 22-25.
20. Shamatov I., Shopulotova Z. ADVANNAGES OF PALATE LASER THERAPY IN COMPLEX TREATMENT OF LARINGITIS //International Bulletin of Medical Sciences and Clinical Research. – 2023. – Т. 3. – №. 9. – С. 104-107.
21. Tilyavova S. SIYDIK TUTOLMASLIK RIVOJLANISHIDA RELAX-2 VA JINSIY GORMONLAR O'RNI //Евразийский журнал медицинских и естественных наук. – 2023. – Т. 3. – №. 2. – С. 149-155.